

LARGE-SCALE MASS POWER SPECTRUM FROM PECULIAR VELOCITIES

I. ZEHAVID

Racah Institute of Physics, The Hebrew University, Jerusalem 91904, Israel

This is a brief progress report on a long-term collaborative project to measure the power spectrum (PS) of mass density fluctuations from the Mark III and the SFI catalogs of peculiar velocities.^{1, 2} The PS is estimated by applying maximum likelihood analysis, using generalized CDM models with and without COBE normalization. The application to both catalogs yields fairly similar results for the PS, and the robust results are presented.

1 Introduction

In the standard picture of cosmology, structure evolved from small density fluctuations that grew by gravitational instability. These initial fluctuations are assumed to have a Gaussian distribution characterized by the PS. On large scales, the fluctuations are linear even at late times and still governed by the initial PS. The PS is thus a useful statistic for large-scale structure, providing constraints on cosmology and theories of structure formation. In recent years, the galaxy PS has been estimated from several redshift surveys.³ In this work, we develop and apply likelihood analysis⁴ in order to estimate the *mass* PS from peculiar velocity catalogs. Two such catalogs are used. One is the Mark III catalog of peculiar velocities,⁵ a compilation of several data sets, consisting of roughly 3000 spiral and elliptical galaxies within a volume of $\sim 80 h^{-1}\text{Mpc}$ around the local group, grouped into ~ 1200 objects. The other is the recently completed SFI catalog,⁶ a homogeneously selected sample of ~ 1300 spiral field galaxies, which complies with well-defined criteria. It is interesting to compare the results of the two catalogs, especially in view of apparent discrepancies in the appearance of the velocity fields.^{7, 8}

2 Method

Given a data set $\mathbf{d}$, the goal is to estimate the most likely model $\mathbf{m}$. Invoking a Bayesian approach, this can be turned to maximizing the likelihood function $\mathcal{L} \equiv \mathcal{P}(\mathbf{d}|\mathbf{m})$, the probability of the data given the model, as a function of the model parameters. Under the assumption that both the underlying velocities and the observational errors are Gaussian random fields, the likelihood function can be written as $\mathcal{L} = [(2\pi)^N \det(R)]^{-1/2} \exp\left(-\frac{1}{2} \sum_{i,j}^N d_i R_{ij}^{-1} d_j\right)$, where $\{d_i\}_{i=1}^N$ is the set of observed peculiar velocities and R is their correlation matrix. R involves the theoretical correlation, calculated in linear theory for each assumed cosmological model, and the estimated covariance of the errors.

The likelihood analysis is performed by choosing some parametric functional form for the PS. For each assumed PS, one can calculate the likelihood function and going over the parameter space find the the PS parameters that provide the




Figure 1: Likelihood analysis results for the flat Λ CDM model with $h = 0.6$. $\ln\mathcal{L}$ contours in the $\Omega - n$ plane are shown for SFI (left panel) and Mark III (middle). The best-fit parameters are marked by ‘s’ and ‘m’ on both, for SFI and Mark III respectively. The right panel shows the corresponding PS for the SFI case (solid line) and for Mark III (dashed). The shaded region is the SFI 90% confidence region. The three dots are the PS calculated from Mark III by Kolatt and Dekel (1997),¹⁰ together with their 1σ error-bar.

maximum likelihood. Confidence levels are estimated by approximating $-2\ln\mathcal{L}$ as a χ^2 distribution with respect to the model parameters. Note that this method, based on peculiar velocities, essentially measures $f(\Omega)^2 P(k)$ and not the mass density PS by itself. Careful testing of the method was done using realistic mock catalogs,⁹ designed to mimic in detail the real catalogs.

We use several models for the PS. One of these is the so-called Γ model, where we vary the amplitude and the shape-parameter Γ . The main analysis is done with a suit of generalized CDM models, normalized by the COBE 4-yr data. These include open models, flat models with a cosmological constant and tilted models with or without a tensor component. The free parameters are then the density parameter Ω , the Hubble parameter h and the power index n . The recovered PS is sensitive to the assumed observational errors, that go as well into R . We extend the method such that also the magnitude of these errors is determined by the likelihood analysis, by adding free parameters that govern a global change of the assumed errors, in addition to modeling the PS. We find, for both catalogs, a good agreement with the original error estimates, thus allowing for a more reliable recovery of the PS.

3 Results

Figure 1 shows, as a typical example, the results for the flat Λ CDM family of models, with a tensor component in the initial fluctuations, when setting $h = 0.6$ and varying Ω and n . The left panel shows the $\ln\mathcal{L}$ contours for the SFI catalog and the middle panel the results for Mark III. As can be seen from the elongated contours, what is determined well is not a specific point but a high likelihood ridge, constraining a degenerate combination of the parameters of the form $\Omega n^{3.7} = 0.59 \pm 0.08$, in this case. The right panel shows the corresponding maximum-likelihood PS for the two catalogs, where the shaded region represents the 90% confidence region obtained from the SFI high-likelihood ridge.

These results are representative for all other PS models we tried. For each

catalog, the different models yield similar best-fit PS, falling well within each others formal uncertainties and agreeing especially well on intermediate scales ($k \sim 0.1 h \text{ Mpc}^{-1}$). The similarity, seen in the figure, of the PS obtained from SFI to that of Mark III is illustrative for the other models as well. This indicates that the peculiar velocities measured by the two data sets, with their respective error estimates, are consistent with arising from the same underlying mass density PS. Note also the agreement with an independent measure of the PS from the Mark III catalog, using the smoothed density field recovered by POTENT (the three dots).¹⁰

The robust result, for both catalogs and all models, is a relatively high PS, with $P(k)\Omega^{1.2} = (4.5 \pm 2.0) \times 10^3 (h^{-1} \text{ Mpc})^3$ at $k = 0.1 h \text{ Mpc}^{-1}$. An extrapolation to smaller scales using the different CDM models gives $\sigma_8\Omega^{0.6} = 0.85 \pm 0.2$. The error-bars are crude, reflecting the 90% formal likelihood uncertainty for each model, the variance among different models and between catalogs. The general constraint of the high likelihood ridges is of the sort $\Omega h_{50}^\mu n^\nu = 0.75 \pm 0.25$, where $\mu = 1.3$ and $\nu = 3.7, 2.0$ for ΛCDM models with and without tensor fluctuations respectively. For open CDM, without tensor fluctuations, the powers are $\mu = 0.9$ and $\nu = 1.4$. For the span of models checked, the PS peak is in the range $0.02 \leq k \leq 0.06 h \text{ Mpc}^{-1}$. The shape parameter of the Γ model is only weakly constrained to $\Gamma = 0.4 \pm 0.2$. We caution, however, that these results are as yet preliminary, and might depend on the accuracy of the error estimates and on the exact impact of non-linearities.²

Acknowledgments

I thank my close collaborators in this work A. Dekel, W. Freudling, Y. Hoffman and S. Zaroubi. In particular, I thank my collaborators from the SFI collaboration, L.N. da Costa, W. Freudling, R. Giovanelli, M. Haynes, S. Salzer and G. Wegner, for the permission to present these preliminary results in advance of publication.

References

1. S. Zaroubi, I. Zehavi, A. Dekel, Y. Hoffman and T. Kolatt, *ApJ* **486**, 21 (1997).
2. W. Freudling, I. Zehavi, L.N. da Costa, A. Dekel, A. Eldar, R. Giovanelli, M.P. Haynes, J.J. Salzer, G. Wegner, and S. Zaroubi, *ApJ* submitted (1998).
3. M.A. Strauss and J.A. Willick, *Phys. Rep.* **261**, 271 (1995).
4. N. Kaiser, *MNRAS* **231**, 149 (1988).
5. J.A. Willick, S. Courteau, S.M. Faber, D. Burstein and A. Dekel, *ApJ* **446**, 12 (1995); J.A. Willick, S. Courteau, S.M. Faber, D. Burstein, A. Dekel and T. Kolatt, *ApJ* **457**, 460 (1996); J.A. Willick, S. Courteau, S.M. Faber, D. Burstein, A. Dekel and M.A. Strauss, *ApJS* **109**, 333 (1997).
6. R. Giovanelli, M.P. Haynes, L.N. da Costa, W. Freudling, J.J. Salzer and G. Wegner, in preparation.
7. L.N. da Costa, W. Freudling, G. Wegner, R. Giovanelli, M.P. Haynes and J.J. Salzer, *ApJ* **468**, L5 (1996).
8. L.N. da Costa, A. Nusser, W. Freudling, R. Giovanelli, M.P. Haynes, J.J. Salzer and G. Wegner, *MNRAS* submitted (1997).

9. T. Kolatt, A. Dekel, G. Ganon and J. Willick, *ApJ* **458**, 419 (1996).
10. T. Kolatt and A. Dekel, *ApJ* **479**, 592 (1997).